

Docket No.: M0025.0293
(PATENT)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of:
Satishchandra P. Patel

Application No.: 10/632,970

Confirmation No.: 4466

Filed: August 4, 2003

Art Unit: 1615

For: PHARMACEUTICAL COMPOSITIONS

Examiner: Not Yet Assigned

CLAIM FOR PRIORITY AND SUBMISSION OF DOCUMENTS

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Dear Sir:

Applicant hereby claims priority under 35 U.S.C. 119 based on the following prior foreign application filed in the following foreign country on the date indicated:

<u>Country</u>	<u>Application No.</u>	<u>Date</u>
United Kingdom	0218002.4	August 2, 2002

In support of this claim, a certified copy of the said original foreign application is filed herewith.

Dated: April 19, 2004

Respectfully submitted,

By Edward A. Meilman

Edward A. Meilman


Registration No.: 24,735

DICKSTEIN SHAPIRO MORIN &
OSHINSKY LLP

1177 Avenue of the Americas
41st Floor

New York, New York 10036-2714
(212) 835-1400

Attorney for Applicant





INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated

P. McHoney
8 April 2004

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference **GBP84783**

2. Patent application number
(The Patent Office will fill in this part)

3 Full name, address and postcode of the or of each applicant (underline all surnames)

Patel, Satishchandra Punambhai
27 Yale Court
Livingston
New Jersey 07039
United States of America

0218002.4

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention **Pharmaceutical compositions**

5. Name of your agent (if you have one)
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Marks & Clerk
57 - 60 Lincolns Inn fields
London WC2A 3LS

Patents ADP number (if you know it)

18001

6.If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application No
(if you know it)

Date of filing
(day / month / year)

7.If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application


Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

NO



ALC
314103
JG

8438194001


~~8438186001~~

2 AUG 2002

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	0
Description	10
Claim(s)	3
Abstract	1
Drawing(s)	



10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77) 1

Request for substantive examination (Patents Form 10/77) 1

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature *Marks & Clerk*

Date: 2 August 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

GB Patent Filings
0207 400 3000

M&C Folio: GBP84783

Document : 776006

Pharmaceutical Compositions

The present invention relates to pharmaceutical compositions, in particular a micro-emulsion concentrate for cyclosporins.

The cyclosporins are a class of cyclic undecapeptides, with important pharmacological activities, in particular immunosuppressive, anti-inflammatory and/or anti-parasitic activities. The first of the cyclosporins to be isolated, and the most commonly known cyclosporin, is Cyclosporin A, formulations of which are commercially available under the trade marks SANDIMMUNE (RTM) and NEORAL (RTM).

The cyclosporins are very lipophilic and hydrophobic compounds, which are sparingly soluble in water, but dissolve readily in organic solvents such as methanol, ethanol, chloroform and the like. The low solubility in water results in extremely low bioavailability of the cyclosporins when administered orally. This may lead to higher dosages being required, with the consequent possibility of undesirable side effects. Therefore, to provide an effective therapeutic concentration of the drug in the body when administered orally represents a considerable challenge. Extensive research has been conducted to find cyclosporin formulations that are effective for oral administration. There are a number of preparations of cyclosporins suitable for oral administration proposed by the prior art.

Prior art formulations of cyclosporins for oral administration have often involved combinations of the cyclosporin with a surfactant, an oil, and a co-surfactant. Such formulations have been intended to be diluted with water prior to drinking. However, this is rather inconvenient, and also the resulting aqueous composition has an unpleasant taste.

In order to alleviate the problems of having to dilute the composition with water prior to oral administration, and the unpleasant taste of the resulting solution, liquid

compositions have been formulated into soft capsule preparations. For example, the formulation commercially available under the trade mark SANDIMMUNE (RTM) is encapsulated in a soft capsule with a gelatine shell. The formulation contains ethanol in order to solubilise the cyclosporin. However, the ethanol can permeate the gelatine shell of the capsule and is volatile at room temperature. This means that the composition of the contents can vary during storage. If too much ethanol is lost, the cyclosporin may precipitate from the composition, with adverse effects on the bioavailability. This results in uncertainties about dosage.

US 4,388,307 discloses compositions comprising a cyclosporin together with at least one of the following components:

- a) a trans-esterification product of a natural or hydrogenated vegetable oil triglyceride and a polyalkylene polyol;
- b) a saturated fatty acid triglyceride; and
- c) a mono- or di-glyceride.

It is preferred that ethanol be used as a further solubilising agent, and the compositions for oral administration disclosed in the Examples all contain ethanol.

US 5,342,625 discloses pharmaceutical compositions comprising cyclosporins in micro-emulsion pre-concentrate and micro-emulsion form. The compositions contain a cyclosporin disposed in a composition comprising a hydrophilic phase, a lipophilic phase and a surfactant. The hydrophilic phase comprises 1,2-propylene glycol or $R_1-(O-(CH_2))_X-OR_2$ wherein R_1 is a C_{1-5} alkyl or a tetrahydrofurfuryl group, R_2 is a C_{1-5} alkyl or a tetrahydrofurfuryl group or is hydrogen, and X is from 1 to 6. The lipophilic phase typically comprises a fatty acid triglyceride. The compositions may contain a C_{1-5} alkanol, such as ethanol, as a co-solvent. However, the compositions disclosed in US 5,342,625 include components which are restricted for pharmaceutical use by several regulatory agencies world-wide, including the FDA, because they are not considered "Generally Recognised As Safe" (GRAS) for oral use.

US 5,759,997 discloses pharmaceutical compositions comprising a cyclosporin, a fatty acid triglyceride, and a glycerol fatty acid partial ester or propylene glycol or sorbitol

complete or partial ester. The compositions may also comprise a viscosity reducer, such as the trans-esterification product of a natural vegetable oil triglyceride and a polyalkylene polyol. Ethanol can also be used, but is less preferred. The compositions may also comprise an emulsifying agent, preferably a tenside having a hydrophilic-lipophilic balance (HLB) of at least 10.

US 6,057,289 discloses pharmaceutical compositions comprising cyclosporin and a carrier comprising

- (a) a cyclosporin solubilising agent consisting essentially of C_6 to C_{22} fatty acids; and
- (b) a water-soluble non-ionic surfactant.

The surfactant should have a hydrophilic-lipophilic balance (HLB) greater than 10, and suitable surfactants include polyoxyethylene products of hydrogenated vegetable oils, polyethoxylated castor oils or polyethoxylated hydrogenated castor oil, polyoxyethylene-sorbitan-fatty acid esters, polyoxyethylene castor oil derivatives and the like. The compositions are for forming microemulsions upon contact with an aqueous medium.

US 5,858,401 discloses compositions that comprise a cyclosporin, a medium chain monoglyceride of C_6 to C_{12} fatty acids, having a monoglyceride content of at least 50%, and at least one surfactant. The surfactant may be, for example, polyglycolised glycerides or ethoxylated glycerides having a molecular weight of PEG between 400 and 2000 and a fatty acid chain length between C_6 to C_{18} . The compositions are for forming microemulsions upon contact with an aqueous medium.

Having regard to the state of the art, it is clear that it is desirable to provide further formulations of cyclosporins suitable for oral administration, and in particular ones which can be formulated in capsules such as soft gelatine capsules, and which are emulsion concentrates (that is, homogeneous solutions which on exposure to water or gastrointestinal fluids form an emulsion having a particle size of less than 5 microns), and preferably microemulsion concentrates, which avoid the use of volatile components

such as ethanol, and which utilise compounds which are Generally Recognised As Safe (GRAS).

There is also a continued need to provide cyclosporin formulations for oral administration which can have high cyclosporin concentrations (thereby reducing the size of capsule required for a given dosage), which exhibit high oral bioavailability, and which are stable (in particular stable against precipitation of the cyclosporin) upon storage. It is also desirable that formulations should have as few components as possible, thereby resulting in ease of manufacture.

The present invention aims to provide cyclosporin compositions which, at least to some extent, satisfy these requirements.

According to the present invention, there is provided a pharmaceutical composition suitable for oral administration in the form of a homogeneous solution which on exposure to water or gastrointestinal fluids forms an emulsion having a particle size of less than 5 microns, the solution comprising:

- (a) a pharmaceutically effective amount of a cyclosporin,
- (b) a carrier medium comprising a mixture of mono- and diesters of propylene glycol with fatty acids having from 8 to 10 carbon atoms or with mixtures of such fatty acids, wherein the monoester makes up not more than 60mol% of the mixture, and
- (c) a non-ionic surfactant having a hydrophilic lipophilic balance (HLB) greater than 10.

The present invention is partly based upon the discovery that the carrier medium as defined in (b) above represents a particularly good solvent medium for cyclosporins, and therefore it is possible to avoid co-solvents such as ethanol, propylene glycol, or the like. The compositions according to the present invention accordingly preferably do not have such co-solvents, and in particular preferably do not contain ethanol.

The compositions according to the present invention preferably do not contain appreciable amounts of water, that is, they are substantially water-free.

The compositions according to the present invention exhibit excellent stability upon storage, and high concentrations of cyclosporins in the compositions can be achieved.

The compositions according to the present invention are homogeneous mixtures which exhibit excellent bioavailability of the cyclosporin *in vivo*.

The cyclosporin is preferably Cyclosporin A. The cyclosporin preferably makes up from 1 to 25% by weight of the composition, more preferably makes up from 5 to 20% by weight of the composition, even more preferably makes up from 10 to 20% by weight of the composition, and most preferably makes up from 15 to 20% by weight of the composition. The cyclosporin is present in the composition of the present invention in pharmaceutically effective amounts. These amounts are well-known in the art. For example, when treating chronic inflammations or provoking an immunosuppressive effect, it is preferred that the daily dose ranges from about 3 mg/kg to about 50 mg/kg.

The carrier medium preferably makes up from 20 to 80% by weight of the composition, more preferably makes up from 35 to 60% by weight of the composition, and most preferably makes up from 40 to 55% by weight of the composition. In the carrier medium, the monoester preferably makes up from 50 to 60mol% of the mixture of mono- and diesters.

In a particularly preferred embodiment, the carrier medium consists of a mixture of mono- and diesters of propylene glycol with capric and caprylic acids. In another particularly preferred embodiment, the carrier medium consists of a mixture of mono- and diesters of propylene glycol with caprylic acid. Suitable products are available commercially as IMWITOR (RTM) 408 and CAPRYOL (RTM) PGMC.

The non-ionic surfactant preferably makes up from 5 to 60% by weight of the composition, more preferably makes up from 20 to 50% by weight of the composition, and most preferably makes up from 30 to 40% by weight of the composition.

The Hydrophilic Lipophilic Balance (HLB) of the non-ionic surfactant is greater than 10, more preferably greater than 12 and most preferably greater than 14.

The non-ionic surfactant must be capable of forming a stable emulsion, preferably a fine emulsion (particle size less than 1 micron), and more preferably a microemulsion, of the composition when it is brought into contact with aqueous fluid, such as in the G.I. tract.

The non-ionic surfactant is preferably selected from the group consisting of: polyoxyethylene products of hydrogenated vegetable oils, polyethoxylated castor oils, polyethoxylated hydrogenated castor oil, polyoxyethylene-sorbitan-fatty acid esters, and polyoxyethylene castor oil derivatives. Particularly preferred surfactants are set out in **Table 1**. Mixtures of these surfactants can also be used.

Trade Name	Description
TWEEN (RTM) 20	Polyoxyethylene (20) sorbitan monolaurate
TWEEN (RTM) 40	Polyoxyethylene (20) sorbitan monopalmitate
TWEEN (RTM) 60	Polyoxyethylene (20) sorbitan monostearate
TWEEN (RTM) 80	Polyoxyethylene (20) sorbitan monooleate
NIKKOL (RTM) HCO30	PEG-30 hydrogenated castor oil
NIKKOL (RTM) HCO40	PEG-40 hydrogenated castor oil
NIKKOL (RTM) HCO50	PEG-50 hydrogenated castor oil
NIKKOL (RTM) HCO60	PEG-60 hydrogenated castor oil
CREMOPHORE (RTM) RH40	Polyoxyethylene 40 castor oil
CREMOPHORE (RTM) RH60	Polyoxyethylene 60 castor oil
CREMOPHORE (RTM) EL35	Polyoxyethylene 35 castor oil

Table 1

The pharmaceutical compositions according to the present invention may further comprise an antioxidant. This antioxidant, when present, is preferably present in an amount of from 0.01% to 2% by weight of the composition, and more preferably from 0.5 to 1% by weight of the composition. The antioxidant may be any suitable

antioxidant, such as are well known to those skilled in the art. Particularly preferred antioxidants are butyl hydroxy anisole (BHA), butyl hydroxy toluene (BHT), and alpha-tocopherol.

Other additives, excipients, and diluents normally used in the pharmaceutical arts may optionally be added to the composition. These include thickening agents, dispersing agents, flavouring agents, sweetening agents, colouring agents, stabilising agents (including pH stabilisers), and preservatives. However, the compositions of the present invention preferably consist only of the components defined in Claim 1, or at least comprise at least 90%, more preferably at least 95%, and more preferably at least 98% by weight of the components defined in Claim 1.

The pharmaceutical compositions according to the present invention may be formulated as a drinking solution, or as a hard or soft capsule. Soft capsule formulations are particularly preferred. Gelatine capsules are also preferred.

The pharmaceutical compositions according to the present invention can be conveniently prepared by uniformly and thoroughly mixing the carrier medium, the cyclosporin, and the surfactant together at room temperature or at slightly elevated temperature, such as a temperature up to 40°C, until a clear solution is obtained, and then cooling the composition to room temperature. The other additives indicated above are then thoroughly admixed therewith. The cyclosporin remains in solution and does not crystallise or precipitate out.

Compositions according to the present invention are preferably for administration to mammals, and especially to humans. It is preferred that the pharmaceutical compositions of the present invention are administered in capsule, liquid-oral, drink solution, or the like form. In a preferred embodiment, the composition is in a form adapted for oral administration in oral unit dosage form. Capsules, e.g., soft or hard gelatine capsules, which represent the preferred oral dosage form, are specially suitable unit dosage forms for oral administration.

Oral unit dosage forms in accordance with the present invention will suitably comprise from 5 to 400 mg and more preferably from 20 to 200 mg, e.g., 25, 50, 100, 125, 150, or 200 mg of cyclosporin. The dosage of the drug and the number of times administered to the patient will vary depending on several factors such as: the age of the patient, the severity of the condition of the patient, and past medical history, and will be a matter to be determined by the attending physician.

When the composition of the present invention is prepared in the form of a soft or hard capsule, the composition may be encapsulated in a gelatine shell which contains any conventional plasticizer. Suitable plasticizers are: glycerine, sorbitol, hexanetriol propylene carbonate, hexane glycol, sorbitans, tetrahydrofuryl alcohol ether, diethylene glycol monoethyl ether, 1,3-trimethyl-2-imidazolidone, dimethylisorbide, and mixtures of these. However, the plasticizer is not limited to those just mentioned, and any suitable plasticizer can be used.

Encapsulation can be achieved by standard techniques which are well known in the art.

Compositions according to the present invention exhibit high solubility of cyclosporin, thereby reducing the size of the capsule or other oral unit dosage form. They also employ only materials that are GRAS for oral use.

The invention will now be further described with reference to the following Examples, it being understood that these are intended to illustrate the invention, and in no way to limit its scope.

Examples

The examples used the ingredients and in the amounts indicated in **Table 2**.

Cyclosporin A was dissolved in the carrier medium, the surfactant was added, and the mixture was mixed for from 10 to 30 minutes at room temperature until the solution was homogeneous.

The solution was then stored overnight up to 24 hours to ensure that no crystallisation occurred.

Ingredients	Example 1	Example 2	Example 3	Example 4
	Weight/mg	weight/mg	weight/mg	weight/mg
Cyclosporin A	100	25	100	100
Mono/diester of propylene glycol with caprylic acid (IMWITOR 408)	285	100	285	170
Mono/diester of propylene glycol with caprylic acid (CAPRYOL PGMC)				130
Polyoxyethylene 35 castor oil (CREMOPHORE EL35)	195		195	180
Polyoxyethylene 20 sorbitan monolaurate (TWEEN 20)		80		
Alpha tocopherol			5	
TOTAL	580 mg	205 mg	585 mg	580 mg

Table 2

The verify that an emulsion was formed, one part of each composition was added to 10 parts of water and stirred gently. There was formed a fine emulsion having a particle size of less than 5 microns, and the Cyclosporin A did not precipitate or crystallise out.

The compositions are suitable for encapsulation into a hard or soft gelatine capsule.

Comparative Examples

The formulation of Example 1 was compared with analogous examples in which the carrier medium was a diester of propylene glycol with capric/caprylic acids (MIGLYOL (RTM) 840) (Comparative Example 1); a mixture of mono- and diesters of propylene glycol with lauric acid (LAUROGLYCOL FCC) (Comparative Example 2); and a

mixture of mono and di-glycerides of capric/caprylic acids (CAPMUL (RTM) MCM) (Comparative Example 3).

The compositions were prepared by mixing the components set out in Table 3 at room temperature until a clear solution was formed. The resulting solutions were maintained at room temperature for four weeks and then compared. The results are shown in Table 4.

Ingredients	Example 1	Comparative Example 1	Comparative Example 2	Comparative Example 3
	weight/mg	weight/mg	weight/mg	weight/mg
Cyclosporin A	100	100	100	100
Mono/diester of propylene glycol with caprylic acid (IMWITOR 408)	285			
Diester of propylene glycol with capric/caprylic acids (MIGLYOL (RTM) 840)		285		
Mono/diester of propylene glycol with lauric acid (LAUROGLYCOL FCC)			285	
Mono/di-glycerides of capric/caprylic acids (CAPMUL(RTM) MCM)				285
Polyoxyethylene 35 castor oil (CREMOPHORE EL35)	195	195	195	195
Alpha tocopherol	5	5	5	5
TOTAL	585 mg	585 mg	585 mg	585 mg

Table 3

Conditions	Example 1	Comparative Example 1	Comparative Example 2	Comparative Example 3
	Observation			
Initial	Clear solution	Hazy solution	Clear solution	Clear solution
4 weeks at 25°C	Clear solution	Precipitation hazy solution	Hazy solution	Hazy solution
4 weeks at 40 °C +75% RH	Clear solution	Precipitation hazy solution	Hazy solution	Hazy solution

Table 4

CLAIMS

1. A pharmaceutical composition suitable for oral administration in the form of a homogeneous solution which on exposure to water or gastrointestinal fluids forms an emulsion having a particle size of less than 5 microns, the solution comprising:
 - (a) a pharmaceutically effective amount of a cyclosporin,
 - (b) a carrier medium comprising a mixture of mono- and diesters of propylene glycol with fatty acids having from 8 to 10 carbon atoms or with mixtures of such fatty acids, wherein the monoester makes up not more than 60mol% of the mixture, and
 - (c) a non-ionic surfactant having a hydrophilic lipophilic balance (HLB) greater than 10.
2. A pharmaceutical composition according to Claim 1 wherein the monoester makes up from 50 to 60mol% of the mixture of mono- and diesters.
3. A pharmaceutical composition according to Claim 1 or Claim 2, wherein the cyclosporin makes up from 1 to 25% by weight of the composition.
4. A pharmaceutical composition according to Claim 3 wherein the cyclosporin makes up from 5 to 20% by weight of the composition.
5. A pharmaceutical composition according to any of Claims 1 to 4, wherein the carrier medium makes up from 20 to 80% by weight of the composition.
6. A pharmaceutical composition according to Claim 5, wherein the carrier medium makes up from 35 to 60% by weight of the composition.

7. A pharmaceutical composition according to any of Claims 1 to 6, wherein the non-ionic surfactant makes up from 5 to 60% by weight of the composition.
8. A pharmaceutical composition according to Claim 7, wherein the non-ionic surfactant makes up from 20 to 50% by weight of the composition.
9. A pharmaceutical composition according to any of Claims 1 to 8, wherein said carrier medium consists of a mixture of mono- and diesters of propylene glycol with capric and caprylic acids.
10. A pharmaceutical composition according to any of Claims 1 to 8, wherein said carrier medium consists of a mixture of mono- and diesters of propylene glycol with caprylic acid.
11. A pharmaceutical composition according to any of Claims 1 to 10, wherein the non-ionic surfactant is selected from the group consisting of: polyoxyethylene products of hydrogenated vegetable oils, polyethoxylated castor oils, polyethoxylated hydrogenated castor oil, polyoxyethylene-sorbitan-fatty acid esters, polyoxyethylene castor oil derivatives, and mixtures thereof.
12. A pharmaceutical composition according to Claim 11, wherein the non-ionic surfactant is selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate, polyoxyethylene (20) sorbitan monopalmitate, polyoxyethylene (20) sorbitan monostearate, polyoxyethylene (20) sorbitan monooleate, PEG-30 hydrogenated castor oil, PEG-40 hydrogenated castor oil, PEG-50 hydrogenated castor oil, PEG-60 hydrogenated castor oil, polyoxyethylene 40 castor oil, polyoxyethylene 60 castor oil, polyoxyethylene 35 castor oil, and mixtures thereof.
13. A pharmaceutical composition according to any of Claims 1 to 12, comprising an antioxidant.

14. A pharmaceutical composition according to Claim 13, wherein the antioxidant is present in an amount of from 0.01% to 2% by weight of the composition.
15. A pharmaceutical composition according to Claim 13 or Claim 14, wherein the antioxidant is selected from the group consisting of BHA, BHT, and alpha-tocopherol.
16. A pharmaceutical composition according to any of Claims 1 to 15, wherein the cyclosporin is Cyclosporin A.
17. A pharmaceutical composition according to any of Claims 1 to 16, formulated as a drinking solution.
18. A pharmaceutical composition according to any of Claims 1 to 16 formulated as a hard or soft capsule.

ABSTRACT**Pharmaceutical Compositions**

The application discloses a pharmaceutical composition suitable for oral administration in the form of a homogeneous solution which on exposure to water or gastrointestinal fluids forms an emulsion having a particle size of less than 5 microns, the solution comprising:

- (a) a pharmaceutically effective amount of a cyclosporin, in particular Cyclosporin A,
- (b) a carrier medium comprising a mixture of mono- and diesters of propylene glycol with fatty acids having from 8 to 10 carbon atoms or with mixtures of such fatty acids, wherein the monoester makes up not more than 60mol% of the mixture, and
- (c) a non-ionic surfactant having a hydrophilic lipophilic balance (HLB) greater than 10.